



Chicken Feed: the KNF way

www.cgnf-hawaii.org
drakew@gmail.com

Master Cho's Wisdom

- Master Cho's Master is Yamaguchi Miyoso
- Concept is LOVE for animals
- Purpose is to glorify God
- Formula for growing strong digestive system in chicks
- Layer feed formula
- Broiler feed formula
- Considerations in Hawai'i





long intestines increase
digestion

Chick Feed Timeline

- First 3 days
 - Brown Rice only
- Day 3-10
 - hard boiled and mashed eggs
 - bamboo leaves
 - food after frenzy
 - brown rice available
- after
 - food





Pile of brown rice for
chicks

placed 24 hours before arrival



amounts of bamboo and
boiled egg to add for 200
broilers



boiled egg acts as
mothers milk for chicks



Chicks have frenzy for boiled eggs. when done, add bamboo and they eat that.



Food trays are put in and
feed given. Enough so gone
by 10am next day

Be careful increasing decreasing
feed amounts

4/20(목). 입식 400마리
 . 수. 칼 (제산제. 균제제)
 . 인. 칼 (비)

《 사료 배합 》

- 분래 (바가리로) 3개
- 쌀겨 1포

- 왕겨 (브레발이로) 10개

- 강수 (바가리로) 1개

- 청미 (") 3개

- 미생물 4번 (") 3개

- 동발 (") 1개

메추리 알 + 바닷물 1ℓ

인. 칼슘 200 ml = 물의 양이 10ℓ 가 되게

《 병아리 음료 》 * 시간이 지나서 각했음

유산균 500 배

한방영양제 700 배

숙(미나리)발효액 300 배

현미 식초 200 배

미네랄 A 1.000 배

chick feed formula

water formula

Feed Theory

- Layers for longevity
 - restricted diet
- Broilers for fast growth
 - as much as they can eat




General formula

- 30% fresh greens
- 30% any fruits, sweet potato, left over food, imo4
- 30% store feed
- 10% IMO5



Feed Mixes

- System 1
 - Protein from eggs
- System 2
 - IMO5 Based
- Hawaii Protein Concerns
 - Hemp seed
 - Maggot system





juicy grasses are good
for feed



feed mixer for making
layer food



Making IMO5

half animal waste; half IMO4



Making IMO6

2 parts IMO4; 1 part IMO5;
yeast and lab water



Thoroughly mixing feed



Fermented 3 days in
box



5 gallon bucket per 120
layers